Local Infusion

MYLOCALINFUSION.COM

COSENTYX (SECUKINUMAB) ORDER FORM

PATIENT INFORMATION

Patient Name: _____ DOB: _____

Mobile Number: _____ Patient Weight: _____kg

Allergies: _____

❑ Patient has known difficult venous access. Comments: _____

❑ New Start Therapy | ❑ Continuation of Therapy & Date of last dose (if applicable): _____

DIAGNOSIS (Provider must specify)

❑ Psoriatic Arthritis, ICD 10: L40.5_____ ❑ Ankylosing Spondylitis, ICD 10: M45.9_____

❑ Non-radiographic Axial Spondyloarthritis, ICD 10: M45.A_____

❑ Other: _____

PROVIDER INFORMATION

Provider Name (print name): _____ Provider NPI: _____

Signature: _____ Date: _____

Contact Name: _____ Phone: _____ Fax: _____

Email Address: _____

Prerequisites to treatment – ensure the following information is complete and attached with referral:

❑ Demographics ❑ Labs and tests supporting diagnosis ❑ Office/progress notes

PRE-MEDICATION (Not typically indicated)

❑ Acetaminophen (Tylenol) 500 mg PO

❑ Benadryl 25mg PO

❑ Famotidine 20 mg IV

❑ Cetirizine (Zyrtec) 10 mg PO

❑ Methylprednisolone (Solu-Medrol) 125 mg IVP

❑ Other: _____ Dose: _____ Route: _____

MEDICATION

MEDICATION	DOSE	ROUTE	FREQUENCY
Cosentyx	❑ With loading dose: 6mg/kg given at week 0, followed by 1.75mg/kg every 4 weeks thereafter ❑ Without loading dose: 1.75mg/kg ***Loading dose optional** ***Max. maintenance dose 300mg per infusion**	IV	q4wks

LABS / SPECIAL INSTRUCTIONS